

October 1979. E. Kiviat. American Museum of Natural History 108367-108369.

Not reported on Jekyll Island by Sandifer *et al.* (1980. Ecological Characterization of the Sea Island Coastal Region of South Carolina and Georgia, Vol. III. Biological Features of the Characterization Area. US Fish Wildl. Serv., Wash. DC. FWS/OBS-79/42 p. 130) or Schwab and Shelton (1981. Bull. Chicago Herpetol. Soc. 16(3):67-69), though known from nearby Little Cumberland and Sapelo Islands.

Submitted by **ERIK KIVIAT**, Hudsonia, Bard College, Annandale, New York 12504. ●

TESTUDINES

CLEMMYS MUHLENBERGI (Bog Turtle) USA: SOUTH CAROLINA: Greenville Co: 2.5 km W Marietta. 23 August 1980. C. E. Putnam, Jr. Verified by D. W. Herman (Atlanta Zoological Park). Clemson University (CU 37-272). One juvenile found basking on a grass clump at 1430 hr; air temperature 25° C. The site is a woodland bog with a canopy of yellow poplar (*Liriodendron tulipifera*) and red maple (*Acer rubrum*). First record for the state and extends the range at least 20 km S from the southern boundary of Transylvania County, North Carolina, where the turtle is known from several localities (Nemuras, K. 1974. Wildl. North Carolina, 38(2):13-15).

Submitted by **DENNIS W. HERMAN**, Department of Herpetology, Atlanta Zoological Park, 800 Cherokee Avenue SE, Atlanta, Georgia 30315 and **CARL E. PUTNAM, JR.**, Rt. 5, Box 37D, Anderson, South Carolina 29621. ●

EMYDOIDEA BLANDINGI (Blanding's Turtle). USA: NEW YORK: Jefferson Co: Mill-site Lake, Town of Theresa. 3 July 1981. R. J. Gawlik. American Museum of Natural History (AMNH 120619). Verified by M. W. Klemens. This report extends the species' northern New York range approximately 10 km to the SE of its reported occurrence at Goose Bay Marsh on the St. Lawrence River (Petokas and Alexander, 1978, SSAR Herp. Review 9(3):107) and more clearly defines its range in the St. Lawrence River lowland as described by Petokas and Alexander (1981, New York Fish and Game J. 28(1):119-120). Mill-site Lake is deep and cold with steep, rocky shorelines having little aquatic vegetation, and thus does not appear to be suitable Blanding's turtle habitat. We suspect that this individual wandered from its more usual habitat such as the ponds and marshes found in the general area of capture.

Submitted by **PETER J. PETOKAS**, U. S. Fish and Wildlife Service, 100 Grange Place, Room 202, Cortland, New York 13045 and **RANDY J. GAWLIK**, Department of Biological Sciences, State University of New York, Cortland, New York 13045. ●

SAURIA

CNEMIDOPHORUS SEXLINEATUS SEXLINEATUS (Six-lined Racerunner). USA: GEORGIA: Glynn Co: Jekyll Island, 1 km S of eastern terminus of Fortson Parkway. 10

October 1979. E. Kiviat. American Museum of Natural History 120073.

Under concrete fragment in low herbs back of ocean beach. Not reported on Jekyll Island by Sandifer *et al.* (1980. Ecological Characterization of the Sea Island Coastal Region of South Carolina and Georgia, Vol. III. Biological Features of the Characterization Area. US Fish Wildl. Serv., Wash. DC. FWS/OBS-79/42 p. 131) or Schwab and Shelton (1981. Bull. Chicago Herpetol. Soc. 16(3):67-69), though known from nearby Little Cumberland and Sapelo Islands.

Submitted by **ERIK KIVIAT**, Hudsonia, Bard College, Annandale, New York 12504. ●

EREMIAS CF. NEUMANNI Tornier (Neumann's Sand-Lizard) AFRICA: TANZANIA: Coast Region, Kisarawe District, Edge of Pugu Forest Reserve, on grounds of Pugu kaolin works, app. 23 km SW Dar es Salaam. 9 January 1971. In short grass on kaolin sand. K. M. Howell, D. M. Pearson. Minaki Secondary School, 23 km SW Dar es Salaam, 10 October 1978. K. M. Howell, Michael O'Meara, Tessa van der Willigen. Specimens identified by Dr. E. N. Arnold, British Museum (Natural History) and catalogued as BMNH 1975.993 and 1978.1331, respectively. These two specimens are the first records of *E. neumanni*-like animals south of Ngatana, Tana River, Kenya (Loveridge, 1957, Bull. Mus. Comp. Zool. 117(2):233).

Submitted by **K. M. HOWELL**, Department of Zoology, P.O. Box 35064, University of Dar es Salaam, Dar es Salaam, Tanzania. ●

EUMECES INEXPECTATUS (Southeastern Five-lined Skink). USA: GEORGIA: Glynn Co: Jekyll Island, pond between Amphitheater and Golf Clubhouse, S of Captain Wylly Road. 12 October 1979. E. Kiviat. American Museum of Natural History 120072.

Under old rubber doormat at wooded edge of pond. Not reported on Jekyll Island by Sandifer *et al.* (1980. Ecological Characterization of the Sea Island Coastal Region of South Carolina and Georgia, Vol. III. Biological Features of the Characterization Area. US Fish Wildl. Serv., Wash. DC. FWS/OBS-79/42 p. 131) or Schwab and Shelton (1981. Bull. Chicago Herpetol. Soc. 16(3):67-69), though known from nearby Cumberland and Sapelo Islands.

Submitted by **ERIK KIVIAT**, Hudsonia, Bard College, Annandale, New York 12504. ●

HEMIDACTYLUS TURCICUS (Mediterranean Gecko). USA: ALABAMA: Jefferson Co: railyard in city limits of Birmingham (T17S, R3W, Sec. 27, N½). 7 October 1981. G. Bosworth. Identified by K. R. Marion. University of Alabama at Birmingham Vertebrate Museum (VM-UAB 0368). First record for county and north Alabama. The locality is ca. 230 km NW of an isolated breeding colony near a motel in Eufaula, Barbour County (Mount, R. H. 1975. The Reptiles and Amphibians of Alabama, Auburn University Press) and ca. 150 km NW of the only other reported locality in

the state, an apparent small breeding population on the Auburn University Campus, Lee County (R. H. Mount, pers. comm.). The specimen was captured under a railroad tie and was evidently introduced by rail from points south. The railyard regularly receives freight shipments from Pensacola, Florida. Several "small lizards" were also seen by workers in the immediate area in the weeks preceding the capture of this specimen, an adult.

Submitted by **KEN R. MARION** and **GREG BOSWORTH**, Biology Department, University of Alabama in Birmingham, University Station, Birmingham, Alabama 35294. ●

SCELOPORUS COUCHI (Couch's Spiny Lizard). MEXICO: NUEVO LEON: Picachos Mts., 9.8 km SW Cerralvo and 10.1 km W on Rancho El Milagro, 11 July 1977, Ernest A. Liner and Allan H. Chaney, EAL 4299 (7 specimens); 9.8 km SW Cerralvo and 10 km W on Rancho El Milagro, Picachos Mts., 15 August 1978, EAL 4560 (4 specimens); 9.8 km SW Cerralvo and 13.3 km W on Rancho El Milagro, Picachos Mts., 15 August 1978, EAL 4564 (1 specimen); 9.8 km SW Cerralvo and 16.7 km W on Rancho El Milagro, Picachos Mts., 16 August 1978, Ernest A. Liner, Richard M. Johnson and Allan H. Chaney, EAL 4569 (4 specimens); verified by Ernest A. Liner. These specimens extend the range eastward approximately 60 km from the nearest known range in the vicinity of Sabinas Hidalgo.

Submitted by **RICHARD M. JOHNSON**, 4170 Lantern Lane, Mobile, Alabama 36609, **ERNEST A. LINER**, 310 Malibou Blvd., Houma, Louisiana 70360 and **ALLAN H. CHANEY**, 1208 W. Richard, Kingsville, Texas 78363. ●

SCELOPORUS POINSETTI POINSETTI (Crevice Spiny Lizard). MEXICO: NUEVO LEON: Picachos Mts., 9.8 km WW Cerralvo and 11.9 km W on Rancho El Milagro, 10 July 1977, EAL 4296 (1 specimen); Picachos Mts., 9.8 km SW Cerralvo and 11.9 km W on Rancho El Milagro, 11 July 1977, EAL 4297 (5 specimens); Picachos Mts., 9.8 km SW Cerralvo and 10.1 km W on Rancho El Milagro, 11 July 1977, Ernest A. Liner and Allan H. Chaney, EAL 4301 (1 specimen); 9.8 km SW Cerralvo and 13.3 km W on Rancho El Milagro, Picachos Mts., 15 August 1978, EAL 4565 (1 specimen); 9.8 km SW Cerralvo and 16.7 km W on Rancho El Milagro, Picachos Mts., 16 August 1978, Ernest A. Liner, Richard M. Johnson and Allan H. Chaney, EAL 4572 (1 specimen); verified by Ernest A. Liner. These records extend the range approximately 60 km east from the previously known range.

Submitted by **ALLAN H. CHANEY**, 1208 W. Richard, Kingsville, Texas 78363, **ERNEST A. LINER**, 310 Malibou Blvd., Houma, Louisiana 70360 and **RICHARD M. JOHNSON**, 4170 Lantern Lane, Mobile, Alabama 36609. ●

SERPENTES

ELAPHE SUBOCULARIS (Trans-Pecos Rat Snake). MEXICO: NUEVO LEON: 35.4 km SW Bustamante, 610 m, 26 July 1979, (fig. 1); 28.3 km W Bustamante, 610 m, 28 July 1981, (fig.